

Index of Symbols

- $(\mathfrak{A}, a)_{a \in A} \models_g \varphi$ The Value of φ in $(\mathfrak{A}, a)_{a \in A}$ is g , 176
 $(\mathfrak{B}, b_a)_{a \in A}$ \mathcal{L}_A -Structure Corresponding to the \mathcal{L} -Structure \mathfrak{B} , 176
 B^Θ Subuniverse Generated by B and Θ , 166
 C (Closure) G -Operator on a G -Algebra, 89
 C Structural Closure Operator, 8
 $C(\Gamma)$ G -Theory Generated by the G -Set of Formulas Γ , 25
 $C \leq C'$ Weaker Than, 9
 C^* Closure G -Operator $C/\tilde{\Omega}(\mathbb{L})$ for a G -Logic $\mathbb{L} = \langle \mathcal{A}, C \rangle$, 101
 $C_{\mathbb{L}}$ Closure G -Operator Induced by a G -Logic, 109
 $C_{\mathcal{A}}$ Closure Operator Induced by the G -Algebra \mathcal{A} , 39
 $C_{\mathbb{K}}$ Closure Operator Induced by the Class \mathbb{K} of G -Algebras, 39
 $C_{\mathbb{K}}$ Equational Consequence Operator Induced by the Class \mathbb{K} of \mathcal{L} -Algebras, 9
 $C_{\mathbb{L}}$ Closure G -Operator Induced by a Class \mathbb{L} of G -Logics, 109
 $C_{\mathbb{M}}$ Closure Operator Induced by the Class \mathbb{M} of G -Matrices, 27
 $C_{\mathbb{M}}$ Consequence Operator Induced by the Class of Matrices \mathbb{M} , 9
 $C_{\mathfrak{A}}$ Closure Operator Induced by the G -Matrix \mathfrak{A} , 27
 $D(\mathfrak{A})$ Diagram of \mathfrak{A} , 176
 $D_\ell(\mathfrak{A})$ Leibniz Diagram of \mathfrak{A} , 176
 $E \text{ comp } F$ G -Congruence E Compatible With G -Filter F , 86
 $E^{\mathcal{E}}$ G -Filter Corresponding to the G -Congruence E , 49
 E^t Translation Associated With the Hybrid Translation E , 45
 $E^{\varphi, \psi}$ Instantiation of E to $\varphi \approx \psi$, 43
 $F^{\mathcal{F}}$ G -Congruence Corresponding to the G -Filter F , 51
 F^t Translation Associated With the Hybrid Translation F , 46
 $F_{\mathbf{A}}^{\delta \approx \varepsilon}$ Filter Generated by the Translation $\delta \approx \varepsilon$ on the Algebra \mathbf{A} , 11
 $H_{\mathbb{K}} : \text{Th}(\mathbf{S}_{\mathbb{K}}) \rightarrow \text{Th}(\mathcal{S})$ Operator Associated with a G -Algebraic Semantics, 58
 $L(\mathfrak{A})$ Infinitary Conjunctions of Leibniz Sentences of $\mathcal{L}_{\mathcal{A}}$, 176
 $[F]_{\Omega}$ \sim_{Ω} -Equivalence Class of F , 130
 $[X]$ Universe of Subalgebra

- Generated by X , 145
- $\Delta(x, y)$ Translation from Equations to Formulas, 10
- $\Delta : \mathcal{S}_K \rightarrow \mathcal{S}$ Interpretation from \mathcal{S}_K to \mathcal{S} , 11
- $\Delta_{\mathbf{A}}$ Least G -Congruence on an G -Algebra \mathbf{A} , 76
- $\Gamma \leq_f \Delta$ $\Gamma \leq \Delta$ and Γ Finite, 26
- $\Gamma \models \Phi$ Semantic Consequence (based on Full Model Class), 171
- $\Gamma \models^* \Phi$ Semantic Consequence (based on Reduced Model Class), 171
- Γ^φ Instantiation of Γ to φ , 43
- Ω Leibniz Operator on the Formula Algebra, 36
- $\Omega(\mathbf{Th}(\mathcal{S}))$ Complete Lattice of the Images of Theories Under the Leibniz Operator, 67
- $\Omega(\mathcal{A})$ Leibniz G -Congruence of \mathfrak{A} , 36
- $\Omega(\mathfrak{A})$ Leibniz G -Congruence of \mathfrak{A} , 153
- $\Omega(\mathfrak{A})$ Leibniz G -Congruence of the G -Matrix \mathfrak{A} , 87
- $\Omega(\mathfrak{A})$ Leibniz Congruence of the Matrix \mathfrak{A} , 10
- $\Omega_{\mathbf{A}}$ Leibniz Operator on the Algebra \mathbf{A} , 10
- $\Omega_{\mathbf{A}}(F)$ Leibniz G -Congruence of F on \mathbf{A} , 36
- $\Omega_{\mathbf{A}}(F)$ Leibniz Congruence of the Matrix $\mathfrak{A} = \langle \mathbf{A}, F \rangle$, 10
- $\Omega_{\mathbf{A}} : G^A \rightarrow \text{Gon}(\mathbf{A})$ Leibniz Operator on \mathbf{A} , 36
- $\Omega_{\mathcal{A}}$ Leibniz Operator of G -Algebra \mathcal{A} , 89
- $\Omega_{\mathcal{A}}(F)$ Leibniz G -Congruence of the G -Matrix $\mathfrak{A} = \langle \mathcal{A}, F \rangle$, 87
- $\Omega_K(\mathbf{Th}(\mathcal{S}))$ Complete Lattice on $\Omega_K(\text{Th}(\mathcal{S}))$ Under the Order Inherited by $\mathbf{Th}(\mathcal{S}_K)$, 60
- $\Omega_K : \text{Th}(\mathcal{S}) \rightarrow \text{Th}(\mathcal{S}_K)$ Operator Associated with a G -Algebraic Semantics, 58
- $\Phi = \{\Phi_g : g \in G\}$ Graded Collection of \mathcal{L} -Formulas, 170
- Θ G -Congruence, 29
- $\Theta_{\mathcal{A}, h}$ G -Kernel Induced by a G -Algebra $\mathcal{A} = \langle \mathbf{A}, E \rangle$ and an $h : \mathbf{Fm}_{\mathcal{L}}(V) \rightarrow \mathbf{A}$, 68
- $\bar{\Theta}$ G -Congruence Induced by Θ , 160
- $\bar{\Theta}$ Reduced G -Congruence Induced by Θ on $\mathbf{A}/\hat{\Theta}$, 76
- $\mathbf{A} = \langle A, F^{\mathfrak{A}} \rangle$ Underlying F -Algebra of \mathcal{L} -Structure \mathfrak{A} , 141
- $\mathbf{A} = \langle A, \mathcal{L}^{\mathbf{A}} \rangle$ \mathcal{L} -Algebra, 9
- \mathbf{A}^* Underlying Algebra of \mathfrak{A}^* , 169
- $\mathbf{Fi}_{\mathcal{S}}(\mathbf{A}) = \langle \text{Fi}_{\mathcal{S}}(\mathbf{A}), \subseteq \rangle$ Complete Lattice of \mathcal{S} -Filters on an Algebra \mathbf{A} , 9
- $\mathbf{Fm}_{\mathcal{L}}(V)$ Absolutely Free Algebra of \mathcal{L} -Formulas, 23
- $\mathbf{Fm}_{\mathcal{L}}(V) = \langle \text{Fm}_{\mathcal{L}}(V), \mathcal{L} \rangle$ Absolutely Free \mathcal{L} -Algebra, 85
- $\mathbf{Gon}(\mathbf{A}) = \langle \text{Gon}(\mathbf{A}), \leq \rangle$ Complete Lattice of G -Congruences on \mathbf{A} , 31, 152
- $\mathbf{Gon}(\mathbb{L})$ Complete Lattice of Logical G -Congruences of \mathbb{L} , 91
- $\mathbf{Gon}(\mathcal{A}) = \langle \text{Gon}(\mathcal{A}), \leq \rangle$ Complete Lattice of G -Congruences on \mathcal{A} , 152
- $\mathbf{G} = \langle G, \leq \rangle$ A Poset of Truth Values With Additional Structure, 23
- $\mathbf{Log}_{\mathbf{G}}(\mathcal{L}) = \langle \text{Log}_{\mathbf{G}}(\mathcal{L}), \leq \rightarrow \rangle$ Complete Lattice of G -Logics over \mathcal{L} , 25
- $\mathbf{Log}_{\mathbf{G}}(\mathcal{A}) = \langle \text{Log}_{\mathbf{G}}(\mathcal{A}), \leq \rangle$ Ordered

- Set of G -Logics on the
 G -Algebra \mathcal{A} , 90
 $\mathbf{Log}_G(\mathcal{A})^L$ Lattice of G -Logics on
 \mathcal{A} Extending L , 95
 $\mathbf{Th}(\mathcal{S})$ Complete Lattice of
 Theories of the Logic \mathcal{S} , 9
 $\mathbf{Th}(\mathcal{S}) = \langle \mathbf{Th}(\mathcal{S}), \leq \rangle$ Complete
 Lattice of G -Theories of
 \mathcal{S} , 25
 $\mathbf{Tm}_{\mathcal{L}}(V)$ Absolutely Free
 \mathcal{L} -Algebra, 142
 $\check{\theta}$ G -Congruence Associated With
 the Stratified Congruence
 θ , 33
 $\delta(x) \approx \varepsilon(x)$ Translation from
 Formulas to Equations, 10
 $\delta \approx \varepsilon : \mathcal{S} \rightarrow \mathcal{S}_K$ Interpretation from
 \mathcal{S} to \mathcal{S}_K , 11
 $\hat{\Theta}$ τ -Stratum of $\hat{\Theta}$, 160
 $\hat{\Theta} = \hat{\Theta}_\tau$ τ -Stratum of the Stratified
 Congruence $\hat{\Theta}$, 76
 $\hat{\Theta} = \{\hat{\Theta}_g : g \in G\}$ Stratified
 Congruence Associated
 With the G -Congruence
 Θ , 32
 $\hat{\Theta} = \{\hat{\Theta}_g\}_{g \in G}$ Stratified Congruence
 Associated With Θ , 160
 $\hat{\Theta}_g$ Congruence Associated With
 the G -Congruence Θ and
 $g \in G$, 31
 $\hat{h} : h^{-1}(\mathfrak{B}) \twoheadrightarrow_s h(\mathfrak{A})$ Reductive
 Morphism Induced by
 $h : \mathfrak{A} \rightarrow \mathfrak{B}$, 149
 $\langle \varphi, \psi \rangle$ \mathcal{L} -Equation, 29
 \leftrightarrow Biconditional, 34
 \mathbb{E} Operator of Expansions, 171
 \mathbb{F} Operator of Filter Extensions,
 171
 \mathbb{H} Operator of Morphic Images,
 171
 L Reduction Operator, 171
 $L(K)$ Class of Tarski Reductions
 of Structures in K , 171
 $\mathbb{O} \leq \mathbb{O}'$ Ordering of Class
 Operators, 171
 $\mathbb{O}\mathbb{O}'$ Composition of Class
 Operators, 171
 \mathbb{O}^* Class Operator $L\mathbb{O}$, 171
 \mathbb{P} Operator of Direct Products,
 171
 \mathbb{R} Operator of Reductions, 171
 \mathbb{S} Operator of Substructures, 171
 L/Θ Quotient G -Logic of L by Θ ,
 97
 $L = \langle \mathcal{A}, C \rangle$ G -Logic on the
 G -Algebra \mathcal{A} , 90
 $L \leq L'$ Weaker Than, 90
 $L^* = L/\tilde{\Omega}(L)$ Reduction of a
 G -Logic L , 101
 $\mathbb{S} = \langle \mathcal{Fm}_{\mathcal{L}}(V), C \rangle$ Sentential
 G -Logic, 103
 $\mathbb{S}^* = \mathbb{S}/\tilde{\Omega}(\mathbb{S})$ Lindenbaum-Tarski
 Quotient of \mathbb{S} , 109
 \mathbb{S}_L Sentential G -Logic Induced by
 the G -Logic L , 110
 \mathbb{S}_L Sentential G -Logic Induced by
 the Class L of G -Logics,
 110
 \mathcal{A}/Θ Quotient G -Algebra of \mathcal{A} by
 Θ , 78
 $\mathcal{A} = \langle \mathbf{A}, E \rangle$ G -Algebra, 38
 $\mathcal{A} = \langle \mathbf{A}, E \rangle$ Underlying G -Algebra
 of \mathcal{L} -Structure \mathfrak{A} , 141
 $\mathcal{A} \leq \mathcal{A}'$ \mathcal{A} a Subalgebra of the
 G -Algebra \mathcal{A}' , 84
 $\mathcal{A} \subseteq_{sd} \prod_{i \in I} \mathcal{A}_i$ \mathcal{A} Subdirect Product
 of the G -Algebras \mathcal{A}_i , 84
 \mathcal{A}^* Algebra $\mathbf{A}/\tilde{\Omega}(L)$ for a G -Logic
 $L = \langle \mathcal{A}, C \rangle$, 101
 \mathcal{A}^* Underlying G -Algebra of \mathfrak{A}^* ,
 169
 $\mathcal{A}^\varepsilon = \langle \mathbf{A}, E^\varepsilon \rangle$ G -Matrix
 Corresponding to
 G -Algebra \mathcal{A} , 49
 \mathcal{C} Collection of Closed G -Sets of
 C , 90
 \mathcal{E} Translation from G -Formulas to
 G -Equations, 41

- $\mathcal{E} : \mathcal{S} \rightarrow \mathcal{S}_K$ Interpretation from a G -Logic \mathcal{S} to a G -2-Logic \mathcal{S}_K , 42
 \mathcal{E}^h Hybrid Translation Associated With the Translation \mathcal{E} , 45
 \mathcal{F} Translation from G -Equations to G -Formulas, 42
 $\mathcal{F} : \mathcal{S}_K \rightarrow \mathcal{S}$ Interpretation from a G -2-Logic \mathcal{S}_K to a G -Logic \mathcal{S} , 42
 $\mathcal{F}^\Theta = \langle \mathbf{Fm}_{\mathcal{L}}(V), \Theta \rangle$ G -Algebra on $\mathbf{Fm}_{\mathcal{L}}(V)$ Induced by the G -Congruence Θ , 68
 \mathcal{F}^h Hybrid Translation Associated With the Translation \mathcal{F} , 46
 $\mathcal{Fm}_{\mathcal{L}}^*(V) = \mathcal{Fm}_{\mathcal{L}}(V)/\tilde{\Omega}(\mathcal{S})$ Lindenbaum-Tarski Algebra of \mathcal{S} , 109
 $\mathcal{Fm}_{\mathcal{L}}(V) = \langle \mathbf{Fm}_{\mathcal{L}}(V), \Delta_{\mathbf{Fm}_{\mathcal{L}}(V)} \rangle$ Absolutely Free G -Algebra, 85
 \mathcal{L} Logical Language, 23
 \mathcal{L}_A , Expansion of \mathcal{L} by Constants for Elements of A , 176
 $\mathcal{S} = \langle \mathcal{L}, C \rangle$ G -2-Logic, 38
 $\mathcal{S} = \langle \mathcal{L}, C \rangle$ G -Logic, 24
 $\mathcal{S} = \langle \mathcal{L}, C \rangle$ Sentential Logic, 8
 $\mathcal{S} = \langle \mathcal{L}, \text{Th}(\mathcal{S}) \rangle$ Sentential Logic, 9
 $\mathcal{S} = \langle \mathcal{L}, \vdash \rangle$ Sentential Logic, 8
 $\mathcal{S} \leq \mathcal{S}'$ \mathcal{S} is a Sublogic of \mathcal{S}' , 24
 $\mathcal{S} \leq \mathcal{S}'$ \mathcal{S}' is an Extension of \mathcal{S} , 24
 $\mathcal{S} \leq \mathcal{S}'$ Weaker Than, 9
 $\mathcal{S}_K = \langle \mathcal{L}, C_K \rangle$ G -2-Logic Induced by the Class K of G -Algebras, 40
 $\mathcal{S}_K = \langle \mathcal{L}, C_K \rangle$ Equational Logic Induced by the Class K of \mathcal{L} -Algebras, 9
 $\mathcal{S}_M = \langle \mathcal{L}, C_M \rangle$ G -Logic Induced by the Class M of G -Matrices, 28
 $\mathcal{S}_M = \langle \mathcal{L}, C_M \rangle$ Logic Induced by the Class of Matrices M , 9
 $\mathcal{Tm}_{\mathcal{L}}(V)$ G -Algebra of \mathcal{L} -Terms, 142
 K^* Class of Tarski Reductions of Structures in K , 171
 K^V Universal Atomic Class Generated by a Class of \mathcal{L} -Structures, 184
 $K^{\mathcal{E}} = \{ \mathcal{A}^{\mathcal{E}} : \mathcal{A} \in K \}$ Class of G -Matrices Corresponding to the Class K of G -Algebras, 49
 K^{V*} Universal Atomic Class Generated by a Class of \mathcal{L} -Structures, 184
 $\text{Log}_{\mathbf{G}}(\mathcal{L})$ Collection of G -Logics over \mathcal{L} , 24
 L^* Collection of Reductions of G -Logics in the Class L , 101
 $\text{Mod}(\Phi)$ Full Model Class of Φ , 170
 $\text{Mod}^*(\Phi)$ Reduced Model Class of Φ , 170
 M^* Class of Reductions of G -Matrices in the Class M , 89
 $M^{\mathcal{F}} = \{ \mathfrak{A}^{\mathcal{F}} : \mathfrak{A} \in M \}$ Class of G -Algebras Corresponding to the Class M of G -Matrices, 51
 \mathfrak{A}/Θ Quotient of \mathfrak{A} by Θ , 162
 $\mathfrak{A}/\Theta = \langle \mathcal{A}/\Theta, \bar{F} \rangle$ Quotient G -Matrix of \mathfrak{A} by Θ , 86
 $\mathfrak{A} = \langle A, F^{\mathfrak{A}}, R^{\mathfrak{A}} \rangle$ \mathcal{L} -Structure, 141
 $\mathfrak{A} = \langle \mathbf{A}, F \rangle$ G -Matrix, 27
 $\mathfrak{A} = \langle \mathbf{A}, F \rangle$ Logical Matrix, 9
 $\mathfrak{A} = \langle \mathcal{A}, F \rangle$ G -Matrix, 86
 $\mathfrak{A} \equiv \mathfrak{B}$ \mathfrak{A} and \mathfrak{B} Elementarily Equivalent, 145
 $\mathfrak{A} \models_X \varphi$ The Universal Closure $\forall \varphi$ Satisfies $\mathfrak{A} \models_X \varphi$, 144
 $\mathfrak{A} \models_X \varphi(x_1, \dots, x_k)[a_1, \dots, a_k]$ For h , such that $h(x_i) = a_i$, for all $1 \leq i \leq k$, $\mathfrak{A} \models \varphi[h]$, 144

- $\mathfrak{A} \models_X \varphi[h]$ For $X \subseteq G$, $\varphi^{\mathfrak{A}}[h] \in X$, 144
 $\mathfrak{A} \lesssim \mathfrak{B}$ \mathfrak{B} a Filter Extension of \mathfrak{A} , 145
 $\mathfrak{A} \subseteq \mathfrak{B}$ \mathfrak{A} a Substructure of \mathfrak{B} , 145
 $\mathfrak{A} \subseteq_e \mathfrak{B}$ \mathfrak{A} Elementary Substructure of \mathfrak{B} , 145
 \mathfrak{A}^* Leibniz Quotient of \mathfrak{A} , 169
 $\mathfrak{A}^* = \mathfrak{A}/\Omega_{\mathfrak{A}}(F)$ Reduction of the G -Matrix \mathfrak{A} , 89
 $\mathfrak{A}^{\mathcal{F}} = \langle \mathbf{A}, F^{\mathcal{F}} \rangle$ G -Algebra Corresponding to G -Matrix \mathfrak{A} , 51
 $\mathfrak{A} \upharpoonright_X$ Substructure of \mathfrak{A} Generated by X , 145
 \mathfrak{B}^{Θ} Substructure Generated by B^{Θ} , 166
 $\nabla_{\mathbf{A}}$ Largest G -Congruence on an G -Algebra \mathbf{A} , 76
 $\overline{\mathbb{P}}$ Class Operator \mathbb{P} Applied with Nonempty Index Sets, 171
 $\pi_{\Theta} : \mathcal{A} \rightarrow \mathcal{A}/\Theta$ Quotient G -Morphism, 78
 $\pi_i : \prod_{i \in I} \mathcal{A}_i \rightarrow \mathcal{A}_i$ i -th Projection G -Morphism, 84
 π_{Θ} Canonical Projection Morphism, 162
 $\prod_{i \in I} \mathcal{A}_i$ Direct Product of G -Algebras, 83
 $\prod_{i \in I} \mathfrak{A}_i$ Direct Product of \mathcal{L} -Structures, 150
 \rightarrow Implication, 34
 $\text{Cl}(C)$ Collection of Closed G -Sets of C , 90
 $\text{Cl}(\mathbb{L})$ Collection of Closed G -Sets of $\mathbb{L} = \langle \mathcal{A}, C \rangle$, 90
 $\text{Gon}(\mathbb{L})$ Collection of Logical G -Congruences of \mathbb{L} , 91
 $\text{Log}_{\mathbf{G}}(\mathcal{A})$ Collection of G -Logics on the G -Algebra \mathcal{A} , 90
 $\theta = \{\theta_g : g \in G\}$ Stratified Congruence, 32
 $\varphi(x_1, \dots, x_n)$ Formula With Free Variables Among x_1, \dots, x_n , 142
 $\varphi \approx \psi$ \mathcal{L} -Equation, 29
 $\varphi^{\mathfrak{A}}[h]$ Truth Value of an \mathcal{L} -Formula φ in a Structure \mathfrak{A} Under an Assignment $h : \mathbf{Tm}_{\mathcal{L}}(V) \rightarrow \mathbf{A}$, 142
 \vdash Structural Consequence Relation, 8
 \widehat{E} Lifting of the G -Set of Equations E , 44
 $\widehat{\Gamma}$ Lifting of the G -Set of Formulas Γ , 44
 $\widetilde{H}_{\mathcal{A}}(\Theta)$ G -logic on \mathcal{A} projectively generated from $\langle \mathcal{A}/\Theta, \text{Fis}_{\mathbb{S}}(\mathcal{A}/\Theta) \rangle$ by $\pi_{\Theta} : \mathcal{A} \rightarrow \mathcal{A}/\Theta$, 120
 $\widetilde{\Omega}(\mathbb{L})$ Tarski G -Congruence of \mathbb{L} , 91
 $\widetilde{\Omega}_{\mathcal{A}}$ Tarski Operator on the G -Algebra \mathcal{A} , 91
 $\widetilde{\Omega}_{\mathcal{A}}(C)$ Tarski G -Congruence of $\mathbb{L} = \langle \mathcal{A}, C \rangle$, 91
 a^* Element in A^* Corresponding to $a \in A$., 169
 e G -Equality Relation Symbol, 145
 $f^{\mathfrak{A}}$ Interpretation of Function Symbol f in the \mathcal{L} -Structure \mathfrak{A} , 141
 $h(\mathfrak{A})$ Image of $\mathfrak{A}' \subseteq \mathfrak{A}$ under $h : \mathfrak{A} \rightarrow \mathfrak{B}$, 147
 $h : \mathbb{L} \cong \mathbb{L}'$ (Logical) G -Isomorphism, 96
 $h : \mathbb{L} \rightarrow \mathbb{L}'$ Logical G -Morphism, 92
 $h : \mathbb{L} \rightarrow_b \mathbb{L}'$ Biological G -Morphism, 93
 $h : \mathcal{A} \rightarrow \mathcal{A}'$ G -Algebra Morphism, 77
 $h : \mathcal{A} \rightarrow \mathcal{A}'$ G -Morphism, 77
 $h : \mathcal{A} \rightarrow \mathcal{B}$ G -Embedding, 146
 $h : \mathfrak{A} \cong \mathfrak{B}$ Isomorphism of \mathcal{L} -Structures, 146

- $h : \mathfrak{A} \rightarrow \mathfrak{B}$ Morphism of \mathcal{L} -Structures, 145
 $h : \mathfrak{A} \rightarrow_e \mathfrak{B}$ Elementary Morphism of \mathcal{L} -Structures, 149
 $h : \mathfrak{A} \rightarrow_s \mathfrak{B}$ Strict Morphism, 146
 $h : \mathfrak{A} \rightarrow \mathfrak{B}$ Embedding of \mathcal{L} -Structures, 146
 $h : \mathfrak{A} \rightarrow_s \mathfrak{B}$ Strict Embedding of \mathcal{L} -Structures, 146
 $h : \mathfrak{A} \rightarrow \mathfrak{B}$ Epimorphism of \mathcal{L} -Structures, 146
 $h : \mathfrak{A} \rightarrow_s \mathfrak{B}$ Reductive Morphism of \mathcal{L} -Structures, 146
 $h^{-1}(\mathfrak{B}')$ Preimage of $\mathfrak{B}' \subseteq \mathfrak{B}$ under $h : \mathfrak{A} \rightarrow \mathfrak{B}$, 147
 $j : \mathcal{A} \hookrightarrow_{sd} \prod_{i \in I} \mathcal{A}_i$ Subdirect Embedding, 84
 $r^{\mathfrak{A}}$ Interpretation of Relation Symbol r in the \mathcal{L} -Structure \mathfrak{A} , 141
 $\mathbf{Con}(\mathfrak{A}) = \langle \mathbf{Con}(\mathfrak{A}), \subseteq \rangle$ Complete Lattice of Congruences of \mathfrak{A} , 10
 $\mathbf{Th}(\mathfrak{S})$ Complete Lattice of \mathfrak{S} -Theories, 105
 $\mathbf{Th}(\mathcal{S}_{\mathfrak{K}}) = \langle \mathbf{Th}(\mathcal{S}_{\mathfrak{K}}), \leq \rangle$ Complete Lattice of G -2-Theories of $\mathfrak{S}_{\mathfrak{K}}$, 41
 \leq Pointwise Ordering of Functions in G^X , 23
 $\mathbf{Alg}(\mathfrak{S})$ Class of \mathfrak{S} -Algebras, 116
 $\mathbf{Alg}^*(\mathfrak{S})$ Class of G -Algebraic Reducts of Reduced \mathfrak{S} -Matrices, 108
 $\mathbf{Atm}(\mathfrak{K})$ Atomic Formulas “Satisfied in \mathfrak{K} ”, 181
 $\mathbf{Cl}(\mathbf{L})^*$ Quotient G -Sets $\mathbf{Cl}(\mathbf{L})/\widetilde{\Omega}(\mathbf{L})$ for a G -Logic $\mathbf{L} = \langle \mathcal{A}, C \rangle$, 101
 $\mathbf{Con}(\mathfrak{A})$ Collection of Congruences of the Matrix \mathfrak{A} , 10
 $\mathbf{Eq}_{\mathcal{L}}(V)$ Set of \mathcal{L} -Equations, 9, 29
 $\mathbf{FMod}(\mathfrak{S})$ Class of Full G -Models of \mathfrak{S} , 113
 $\mathbf{FMod}^*(\mathfrak{S})$ Class of Reduced Full G -Models of \mathfrak{S} , 113
 $\mathbf{FMod}_{\mathfrak{S}}(\mathcal{A})$ Class of Full G -Models of \mathfrak{S} on \mathcal{A} , 113
 $\mathbf{Fi}_{\mathfrak{S}}^{\star}(\mathcal{A})$ Collection of Leibniz \mathfrak{S} -Filters, 129
 $\mathbf{Fi}_{\mathfrak{S}}(\mathcal{A})$ Collection of \mathfrak{S} -Filters on \mathcal{A} , 104
 $\mathbf{Fi}_{\mathfrak{S}}(\mathbf{A})$ Collection of \mathfrak{S} -Filters on \mathbf{A} , 28
 $\mathbf{Fi}_{\mathfrak{S}}(\mathbf{A})$ Collection of \mathfrak{S} -Filters on an Algebra \mathbf{A} , 9
 $\mathbf{Fm}_{\mathcal{L}}(V)$ Collection of \mathcal{L} -Formulas, 140
 $\mathbf{Fm}_{\mathcal{L}}(V)$ Set of \mathcal{L} -Formulas, 23
 $\mathbf{Gon}(\mathbf{A})$ Collection of G -Congruences on \mathbf{A} , 30, 151
 $\mathbf{Gon}(\mathcal{A})$ Collection of G -Congruences on \mathcal{A} , 152
 $\mathbf{Gon}(\mathcal{A})$ Collection of G -Congruences on the (Reduced) G -Algebra \mathcal{A} , 78
 $\mathbf{Gon}(\mathfrak{A})$ Collection of G -Congruences of a G -Matrix \mathfrak{A} , 86
 $\mathbf{Gon}(\mathfrak{A})$ Collection of G -Congruences on \mathfrak{A} , 34, 152
 $\mathbf{Gon}_{\mathfrak{K}}(\mathcal{A})$ Collection of all \mathfrak{K} - G -Congruences on the G -Algebra \mathcal{A} , 85
 $\mathbf{Ker}(h)$ G -Kernel of $h : \mathbf{A} \rightarrow \mathbf{B}$, 154
 $\mathbf{Mat}(\mathfrak{S})$ Collection of \mathfrak{S} -Matrices, 104
 $\mathbf{Mat}^*(\mathfrak{S})$ Collection of Reductions of \mathfrak{S} -Matrices, 104
 $\mathbf{Mod}(\mathfrak{S})$ Collection of G -Models of \mathfrak{S} , 111
 $\mathbf{Th}(\mathfrak{S})$ Collection of \mathfrak{S} -Theories, 105

$\text{Th}(\mathcal{S})$ Collection of G -Theories of \mathcal{S} , 25	G -2-Theories of \mathcal{S}_κ , 40
$\text{Th}(\mathcal{S})$ Set of Theories of the Logic \mathcal{S} , 9	$\text{Tm}_{\mathcal{L}}(V)$ Collection of \mathcal{L} -Terms, 140
$\text{Th}(\mathcal{S}_\kappa)$ Collection of	\sim_Ω Relation on $\text{Fis}(\mathcal{A})$, 129